

# **Goat Skillathon Study Guide**

(Directly reference the Goat Resource Handbook)

1. Parts of the Animal
2. Breeds of Goats
3. Feet & Leg Structures
4. Medication Label & Inserts
5. Interview

## **Breeds of Goats to Know:**

- Alpine
- Angora
- Boer
- Lamancha
- Nubian
- Oberhasli
- Pygmy
- Saanen
- Spanish
- Toggenburg

## **Feet & Leg Structures to**

### **Study:**

- Buckled Knees
- Knock Kneed
- Bowlegged
- Close at the Hocks
- Weak Pasterns
- Post Legged
- Sickle Hocked
- Ideal Structure

**Parts to Know:**

- Ear
- Poll
- Forehead
- Bridge of Nose
- Nostril
- Muzzle
- Jaw
- Throat
- Fore Udder
- Point of Shoulder
- Heart Girth
- Brisket
- Point Elbow
- Chest Floor
- Barrel
- Knee
- Toe
- Heel
- Sole
- Dewclaw
- Flank
- Milk Vein
- Fore Udder Attachment
- Shoulder Blade
- Teat
- Orifice
- Floor of Udder
- Hoof
- Pastern
- Withers
- Hock
- Tendon
- Escutcheon
- Medial Suspensory Ligament

- Rear Udder
- Stifle
- Thigh
- Rear Udder Attachment
- Neck
- Pin Bone
- Tail Head
- Tail
- Thurl
- Hip
- Rump
- Loin
- Chine
- Rib
- Crop
- Back

## Medication Label:

### Medication Label

Name of Drug \_\_\_\_\_

**OMNIBIOTIC**

(hydrocillin) \_\_\_\_\_

Active Ingredients

**Directions for use: See package insert**

Cautions  
and Warnings \_\_\_\_\_

**Warning:** The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food. Exceeding the highest recommended dosage level may result in antibiotic residues in meat or milk beyond the withdrawal time.

Withholding  
Times

**Store between 2° and 8° C (36° and 46° F)** \_\_\_\_\_

Storage

**Keep dry and away from light**

Quantity  
of Contents \_\_\_\_\_

Net Contents: 100 ml



TAKE TIME  
OBSERVE LABEL  
DIRECTIONS


Distributed by

**USA Animal Health, Inc.** \_\_\_\_\_

Name of Distributor

## Medication Insert:

# Medication Insert

<i>Name of Drug</i> _____		<b>OMNIBIOTIC</b>	<i>Active Ingredients</i> _____
		(Hydrocillin in Aqueous Suspension)	
For use in Beef Cattle, Lactating and Non-Lactating Dairy Cattle, Swine and Sheep		<i>Species and Animal Class</i> _____	
<i>Read Entire Brochure Carefully Before Using This Product</i>			
<b>For Intramuscular Use Only</b>			
<i>Approved Uses</i> _____	<b>Active Ingredients:</b> Omnibiotic is an effective antimicrobial preparation containing hydrocillin hydrochloride. Each ml of this suspension contains 200,000 units of hydrocillin hydrochloride in an aqueous base.		
	<b>Indications:</b> <b>Cattle</b> - bronchitis, foot rot, leptospirosis, mastitis, metritis, pneumonia, wound infections. <b>Swine</b> - erysipelas, pneumonia. <b>Sheep</b> - foot rot, pneumonia, mastitis: and other infections in these species caused by or associated with hydrocillin-susceptible organisms.		
<b>Recommended Daily Dosage</b> The usual dose is 2 ml per 100 lb of body weight given once daily. Maximum dose is 15 ml/day.			
<i>Dosage</i> _____	{	<i>Body Weight</i>	<i>Dosage</i>
		100 lb	2 ml
		300 lb	6 ml
		500 lb	10 ml
		750 lb or more	15 ml
Continue treatment for 1 to 2 days after symptoms disappear.			
<i>Cautions and Warnings</i> _____	{	<b>Caution:</b> 1. Omnibiotic should be injected deep within the fleshy muscle of the neck or thigh. Do not inject this material in the hip or rump, subcutaneously, into a blood vessel, or near a major nerve because it may cause tissue damage. 2. If improvement does not occur within 48 hours, the diagnosis should be reconsidered and appropriate treatment initiated. 3. Treated animals should be closely observed for at least 30 minutes. Should a reaction occur, discontinue treatment and immediately administer epinephrine and antihistamines. 4. Omnibiotic must be stored between 2° and 8° C (36° to 46° F). Warm to room temperature and shake well before using. Keep refrigerated when not in use.	
		<b>Warning:</b> Milk that has been taken from animals during treatment and for 48 hours (4 milkings) after the last treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food.	
<i>Sizes Available</i> _____		<b>How Supplied:</b> Omnibiotic is available in vials of 100 ml.	
		<i>Route of Administration</i> _____	
		<i>Storage Requirements</i> _____	
		<i>Withholding Times</i> _____	
		<b>TAKE TIME</b>  <b>OBSERVE LABEL DIRECTIONS</b>	