

Horse Skillathon Study Guide

(Directly reference the Horse Resource Handbook)

1. Parts of the Animal
2. Breeds of Horses
3. Conformation - Front View, Rear View, Front Side View, & Rear Side View
4. Medication Label & Inserts
5. Interview

Parts to Know:

- Poll
- Neck
- Crest
- Shoulder
- Withers
- Back
- Loin
- Point of Hip
- Croup
- Thigh Area
- Buttock
- Flank
- Stifle
- Gaskin
- Hock
- Fetlock Joint
- Coronet
- Pastern
- Ankle
- Cannon
- Heart Girth
- Barrel Area
- Abdomen
- Hoof
- Knee
- Forearm
- Elbow
- Arm
- Chest
- Point of Shoulder
- Throat Latch
- Upper Lip
- Lower Lip
- Muzzle
- Under Lip
- Nostril
- Bridge of Nose
- Face
- Forehead
- Chestnut
- Ergot


Breeds to Study:

- American Paint
- American Saddlebred
- Appaloosa
- Arabian
- Belgian
- Clydesdale
- Donkey
- Miniature Horse
- Morgan
- Mule
- Palomino
- Percheron
- Pony of the Americas
(POA)
- Peruvian Paso
- Quarter Horse
- Shetland Pony
- Shire
- Standardbred
- Suffolk
- Tennessee Walking
Horse
- Thoroughbred
- Welsh Pony

Conformation


- Bow Legged
- Bench Kneed
- Base Narrow
- Knock Kneed
- Pigeon Toed
- Narrow Chested, Toes Out
- Cow Hocked
- Stands Close
- Stands Wide
- Camped Out
- Camped Under
- Calf Kneed
- Buck Kneed
- Sickle Hocked
- Post Legged
- Ideal Position

Medication Label:

Medication Label		
Name of Drug	OMNIBIOTIC	
	(hydrocillin)	Active Ingredients
Directions for use: See package insert		
Cautions and Warnings	Warning: The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food. Exceeding the highest recommended dosage level may result in antibiotic residues in meat or milk beyond the withdrawal time.	Withholding Times
Store between 2° and 8° C (36° and 46° F)		Storage
Keep dry and away from light		
Quantity of Contents	Net Contents: 100 ml	
TAKE TIME	Distributed by	
	Name of Distributor	
OBSERVE LABEL DIRECTIONS	USA Animal Health, Inc.	

Medication Insert:

Medication Insert

<i>Name of Drug</i> _____		OMNIBIOTIC	<i>Active Ingredients</i> _____
		(Hydrocillin in Aqueous Suspension)	
For use in Beef Cattle, Lactating and Non-Lactating Dairy Cattle, Swine and Sheep		<i>Species and Animal Class</i> _____	
<i>Read Entire Brochure Carefully Before Using This Product</i>			
For Intramuscular Use Only			
<i>Approved Uses</i> _____	Active Ingredients: Omnibiotic is an effective antimicrobial preparation containing hydrocillin hydrochloride. Each ml of this suspension contains 200,000 units of hydrocillin hydrochloride in an aqueous base.		
	Indications: Cattle - bronchitis, foot rot, leptospirosis, mastitis, metritis, pneumonia, wound infections. Swine - erysipelas, pneumonia. Sheep - foot rot, pneumonia, mastitis: and other infections in these species caused by or associated with hydrocillin-susceptible organisms.		
Recommended Daily Dosage			
<i>The usual dose is 2 ml per 100 lb of body weight given once daily. Maximum dose is 15 ml/day.</i>			
<i>Dosage</i> _____	{	<i>Body Weight</i>	<i>Dosage</i>
		100 lb	2 ml
		300 lb	6 ml
		500 lb	10 ml
		750 lb or more	15 ml
<i>Continue treatment for 1 to 2 days after symptoms disappear.</i>			
<i>Cautions and Warnings</i> _____	Caution: 1. Omnibiotic should be injected deep within the fleshy muscle of the neck or thigh. Do not inject this material in the hip or rump, subcutaneously, into a blood vessel, or near a major nerve because it may cause tissue damage. 2. If improvement does not occur within 48 hours, the diagnosis should be reconsidered and appropriate treatment initiated. 3. Treated animals should be closely observed for at least 30 minutes. Should a reaction occur, discontinue treatment and immediately administer epinephrine and antihistamines. 4. Omnibiotic must be stored between 2° and 8° C (36° to 46° F). Warm to room temperature and shake well before using. Keep refrigerated when not in use.		<i>Route of Administration</i> _____
	Warning: Milk that has been taken from animals during treatment and for 48 hours (4 milkings) after the last treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food.		<i>Storage Requirements</i> _____ <i>Withholding Times</i> _____
<i>Sizes Available</i> _____	How Supplied: Omnibiotic is available in vials of 100 ml.		TAKE TIME  OBSERVE LABEL DIRECTIONS

